

# Atomic Cross Section for Plasma-wall and Plasma Edge Interaction Modelling in Controlled Fusion Reactor

Saed J. Al Atawneh<sup>1,2</sup> and K Tőkési<sup>1</sup>

<sup>1</sup>Institute for Nuclear Research (ATOMKI), 4026 Debrecen Bem tér 18/c, Hungary

<sup>2</sup> Doctoral School of Physics, Faculty of Science and Technology, University of Debrecen, P.O.400, Debrecen, Hungary

The hydrogen atoms generated by ion recombination on the fusion reactor's wall and other plasma-facing components [1-5] play a significant role in the beam emission spectroscopy (BES), which is an active plasma diagnostic tool used for density measurements in fusion research [1]. The cross sectional data of H-H atom collisions act as raw data for BES modeling. Besides hydrogen atoms, the impurities such as carbon, lithium, oxygen, deuterium, and tritium have existed in the plasma edge area [4, 5]. As comprehensive studies, the interactions of carbon and lithium ions with neutral hydrogen atoms were also investigated [4, 5], where carbon composites are used in first wall tiles [6-8] and lithium ions were used as a potential solution to solve the fusion reactor divertor heat flux. The ionized lithium ions can form a highly radiative layer of plasma, thus could significantly reduce the heat flux to the divertor surfaces [4, 9]. This work is created toward developing a theoretical description of inelastic interactions such as ionization and excitation processes that can give the total cross section accurately. In addition it is also contribute to creating a database for total cross sections as raw data for the BES modeling [10].

Classical calculations for determining atomic collision cross sections have received a great deal of interest in the past 20 years. There was a great revival of the CTMC calculations applied in atomic collisions involving three or more particles [2-5]. The CTMC method is a non-perturbative method, where classical equations of motions are solved numerically [2-5]. For a better description of the classical atomic collisions, the quasi-classical trajectory Monte Carlo model of the Kirschbaum and Wilets (QCTMC) improves the results of the standard CTMC model [3-5].

We present ionization, excitation, and electron capture cross sections in collisions between hydrogen atom, carbon and lithium ions with hydrogen atoms. We found a reasonably good agreement between classical results and the previously obtained experimental data.

## References

- [1] Atawneh S. J. A., Asztalos Ö., Szondy B., Pokol G. I. and Tőkési K. 2020 *Atoms* 8 31
- [2] Saed J.A.A. and Tőkési K. 2021 *J. Phys. B: At. Mol. Opt. Phys.* 54 065202
- [3] Saed J.A.A. and Tőkési K. Ionization cross sections in collisions between two hydrogen atoms by a quasi-classical trajectory Monte Carlo model, (To be submitted)
- [4] Saed J.A.A. and Tőkési K. Target electron removal in  $C^{5+} + H$  collision, (Accepted in Nuclear Fusion)
- [5] Saed J.A.A. and Tőkési K. Ionization Cross Section in  $Li^{9+} + H$  Collision. (To be published in Physics letters A)
- [6] Lawson K.D., Coffey I.H., Aggarwal K.M., Keenan F.P. and Contributors J-E. 2013 *J. Phys. B: At. Mol. Opt. Phys* 46 035701
- [7] Federici G. 2006 *Phys. Scr* T124 1-8
- [8] Lipschultz B., et al 2007 *Nucl. Fusion* 47 1189-1205
- [9] Rognlien T.D. and Rensink M.E. 2002 *Physics of Plasmas* 9 2120-2126
- [10] Saed J.A.A. and Tőkési K. PROJECTILE IONIZATION, EXCITATION AND DE-EXCITATION CROSS SECTION DATABASE IN COLLISION BETWEEN TWO HYDROGEN ATOMS IN THE IMPACT ENERGY RANGE BETWEEN 5.0 keV – 5.0 MeV – PART I, TARGET IS IN GROUND STATE. (To be published in Atomic Data and Nuclear Data tables)